

# Impacts of Lake Erie Harmful Algal Blooms on the Abundance and Growth of Larval Fishes and their Prey Resources

Tomena K. Scholze

William W. Taylor




Edward F. Roseman

Nathan Williams







# Harmful Algal Blooms (HABs)



- HABs are a growing concern worldwide
- Link terrestrial and aquatic ecosystems
- Cultural eutrophication affects water quality and fisheries

# Lake Erie HABs




- 1960s-1970s: cyanobacterial blooms
- 1972: Clean Water Act, Great Lakes Water Quality Agreement
- Recent increase in frequency and intensity
  - 1995-2001: minimal HABs
  - 2002-2007: yearly moderate HABs
  - 2008-2015: yearly severe HABs
- Overlap with critical fish habitat in the western basin




# Research Hypothesis

- H: Lake Erie *Microcystis* blooms will decrease fishery recruitment indirectly by limiting the availability of suitable prey.
  - Specific focus on walleye, a keystone predator that supports a valuable fishery
  - Prey: Clupeids (alewife, gizzard shad), *Notropis* spp. (shiners), rainbow smelt
  - Linked to zooplankton indirectly through these prey fishes
  - Trophic cascade framework



# Trends in YOY CPUE

- Fishery-independent bottom trawl survey to estimate year class strength
- Total CPUE was not significantly different HAB vs. non-HAB years




Produced by Tomena Scholze under written agreement with the Lake Erie Committee (2015).



# Trends in YOY CPUE


- HABs associated with lower clupeid and walleye CPUE



Produced by Tomena Scholze under written agreement with the Lake Erie Committee (2015).

# Trends in YOY CPUE

- Relationship between shiners and HABs depended on species




Produced by Tomena Scholze under written agreement with the Lake Erie Committee (2015).



# Trends in YOY CPUE

- Rainbow smelt CPUE was higher during HAB years




Produced by Tomena Scholze under written agreement with the Lake Erie Committee (2015).



# Trends in YOY CPUE

1995-2001: Minimal, 2002-2007: Moderate, 2008-2015: Severe



- Alewife, spottail shiner CPUE decreased with increasing HAB severity



# Trends in YOY CPUE

1995-2001: Minimal, 2002-2007: Moderate, 2008-2015: Severe

- Gizzard shad, walleye CPUE is highest during minimal HAB years




Produced by Tomena Scholze under written agreement with the Lake Erie Committee (2015).



# Trends in YOY CPUE

1995-2001: Minimal, 2002-2007: Moderate, 2008-2015: Severe

- Rainbow smelt CPUE is lowest during minimal HAB years





Produced by Tomena Scholze under written agreement with the Lake Erie Committee (2015).

# Trends in YOY CPUE

1995-2001: Minimal, 2002-2007: Moderate, 2008-2015: Severe

- Total CPUE, emerald shiner, mimic shiner is highest during moderate HABs




Produced by Tomena Scholze under written agreement with the Lake Erie Committee (2015).



# Trends in YOY CPUE

- Walleye may withstand moderate HABs, if they switch to less desirable prey (Clupeids → Notropis), and if there aren't too many consecutive HABs years




# Trends in YOY length

- Length as a proxy for growth
- Alewife and gizzard shad were significantly longer in non-HAB years
- Emerald shiner, spottail shiner, rainbow smelt were longer in HAB years
- No significant difference in walleye length in HAB vs non-HAB years
- We see a decrease in YOY abundance (year class strength) because of decreased growth and survival during the larval stage




# Trends in larval CPUE and length


- 2015 field study
- Total CPUE, gizzard shad, Notropis spp., and walleye CPUE were higher at sites without HABs
- No significant difference in walleye length at HAB vs non-HAB sites



# Zooplankton Trends




- Zooplankton density is higher at HAB sites
  - Copepod density was higher, no significant difference in Daphnia density
- Suggests larvae are not prey-limited during HABs





# Summary

- Walleye YOY CPUE is lower during any HAB event because their preferred prey (clupeids) are less abundant
- Other prey items (Notropis) are available until HABs are consistently severe
- Larval CPUE (overall) is lower during HABs



# Literature Cited

- Ger et al. 2016. Effect of the toxin (microcystin) content of *Microcystis* on copepod grazing. Harmful Algae 52:34-45.
- International Joint Commission, 2014. A balanced diet for Lake Erie: Reducing phosphorus loadings and harmful algal blooms. Report of the Lake Erie Ecosystem Priority. 96pp.
- Madenjian et al. 1996. First-year growth, recruitment, and maturity of walleyes in western Lake Erie. Trans. Am. Fish. Soc. 125:821-830.
- Michalak et al. 2012. Record-setting algal bloom in Lake Erie caused by agricultural and meteorological trends consistent with expected future conditions. PNAS 110(16): 6448-6452.
- NOAA. 2015. Experimental Lake Erie Harmful Algal Bloom Bulletin. No. 27. November 10, 2015.
- Taylor, W.W et al. 2013. Great Lakes fishery policy and management. A binational perspective. 2<sup>nd</sup> Edition. MSU Press, East Lansing, MI. 865pp.



# Acknowledgements



- R Core Team (2014). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>
- YOY trends: Produced by Tomena Scholze under written agreement with the Lake Erie Committee (2015).
- Wynne, T.T. and R.P. Stumpf. 2015. Spatial and temporal patterns in the seasonal distribution of toxic cyanobacteria in western Lake Erie from 2002-2004. *Toxins* 7(5):1649-1663.

# Questions?

